

## Chapter 5

### Page 265, Practice Problems 5.1.1

- $\text{H}_2\text{SO}_4$  acid
  - $\text{XeF}_6$  molecular
  - $\text{CH}_3\text{COOH}$  acid
  - $\text{NaCH}_3\text{COO}$  salt
  - $\text{KOH}$  base
  - $\text{NH}_3$  molecular (but also a base)
- $\text{CH}_3\text{COOH} + \text{LiOH} \rightarrow \text{LiCH}_3\text{COO} + \text{H}_2\text{O}$   
salt
  - $2\text{HI} + \text{Ca}(\text{OH})_2 \rightarrow 2\text{H}_2\text{O} + \text{CaI}_2$   
salt
  - $3\text{Mg}(\text{OH})_2 + 2\text{H}_3\text{PO}_4 \rightarrow 6\text{H}_2\text{O} + \text{Mg}_3(\text{PO}_4)_2$   
salt

	Parent Acid	Parent Base
3. a) $\text{KNO}_2$	$\text{HNO}_2$	$\text{KOH}$
b) $\text{NH}_4\text{Cl}$	$\text{HCl}$	$\text{NH}_4\text{OH}$ (or $\text{NH}_3$ )
c) $\text{CuC}_2\text{O}_4$	$\text{H}_2\text{C}_2\text{O}_4$	$\text{Cu}(\text{OH})_2$
d) $\text{NaCH}_3\text{COO}$	$\text{CH}_3\text{COOH}$	$\text{NaOH}$

### Page 268, Practice Problems 5.1.2

- $\text{HIO}_3 + \text{NO}_2^- \leftrightarrow \text{HNO}_2 + \text{IO}_3^-$   
acid base acid base
  - $\text{HF} + \text{HC}_2\text{O}_4^- \leftrightarrow \text{H}_2\text{C}_2\text{O}_4 + \text{F}^-$   
acid base acid base
  - $\text{Al}(\text{H}_2\text{O})_6^{3+} + \text{SO}_3^{2-} \leftrightarrow \text{HSO}_3^- + \text{Al}(\text{H}_2\text{O})_5\text{OH}^{2+}$   
acid base acid base
- 

Conjugate acid	Conjugate base
$\text{H}_2\text{O}_2$	$\text{HO}_2^-$
$\text{H}_3\text{BO}_3$	$\text{H}_2\text{BO}_3^-$
$\text{HCOOH}$	$\text{HCOO}^-$
$\text{HC}_6\text{H}_5\text{O}_7^{2-}$	$\text{C}_6\text{H}_5\text{O}_7^{3-}$


- $\text{HNO}_2 + \text{NH}_3 \leftrightarrow \text{NO}_2^- + \text{NH}_4^+$   
acid base base acid  
conjugate pairs:  $\text{HNO}_2/\text{NO}_2^-$  and  $\text{NH}_3/\text{NH}_4^+$
  - $\text{H}_3\text{C}_6\text{H}_5\text{O}_7 + \text{CN}^- \leftrightarrow \text{H}_2\text{C}_6\text{H}_5\text{O}_7^- + \text{HCN}$   
acid base base acid  
conjugate pairs:  $\text{H}_3\text{C}_6\text{H}_5\text{O}_7/\text{H}_2\text{C}_6\text{H}_5\text{O}_7^-$  and  $\text{CN}^-/\text{HCN}$
  - $\text{PO}_4^{3-} + \text{H}_2\text{S} \leftrightarrow \text{HPO}_4^{2-} + \text{HS}^-$   
base acid acid base  
conjugate pairs:  $\text{HPO}_4^{2-}/\text{PO}_4^{3-}$  and  $\text{H}_2\text{S}/\text{HS}^-$

### Page 268, Quick Check

- $\text{HCO}_3^- + \text{CN}^- \leftrightarrow \text{CO}_3^{2-} + \text{HCN}$
- $\text{HCO}_3^- + \text{H}_2\text{O} \leftrightarrow \text{H}_2\text{CO}_3 + \text{OH}^-$
- b) and e) are amphiprotic




### Page 270, 5.1 Review Questions

- Both Arrhenius and Bronsted Lowry acids contain  $\text{H}^+$  ions. Definitions of a base differ. Arrhenius base contains  $\text{OH}^-$  ions. Bronsted Lowry base accepts  $\text{H}^+$  ion.
- A Hydrogen atom contains 1 proton and 1 electron. When a  $\text{H}^+$  ion is formed, the electron is given away, leaving a single proton.
- 




One of the lone pairs of electrons on the oxygen atom of water will attract the  $\text{H}^+$  ion from an acid.


4.




5.



6.



7.



b) HClO

8. a) Add a H<sup>+</sup> ion. Example: CN<sup>-</sup> becomes HCN  
 b) Remove a H<sup>+</sup> ion. Example: HNO<sub>2</sub> becomes NO<sub>2</sub><sup>-</sup>

9. a) HO<sub>2</sub><sup>-</sup>

b) N<sub>2</sub>H<sub>5</sub><sup>+</sup>

c) OC<sub>6</sub>H<sub>5</sub><sup>-</sup>


d) C<sub>6</sub>H<sub>5</sub>NH<sub>3</sub><sup>+</sup>

10. A substance that is able act as both an acid or a base depending on the other substance present. Examples: HSO<sub>3</sub><sup>-</sup>, H<sub>2</sub>PO<sub>4</sub><sup>-</sup>, HPO<sub>4</sub><sup>2-</sup>, H<sub>2</sub>O

11. a) As an acid: HCO<sub>3</sub><sup>-</sup> + H<sub>2</sub>O ↔ CO<sub>3</sub><sup>2-</sup> + H<sub>3</sub>O<sup>+</sup>; As a base: HCO<sub>3</sub><sup>-</sup> + H<sub>2</sub>O ↔ H<sub>2</sub>CO<sub>3</sub> + OH<sup>-</sup>  
 Use litmus paper to test the solution. If it is acidic, the litmus will turn red. If it is basic, the litmus will turn blue.

b) As a base. H<sub>2</sub>O and CO<sub>2</sub> can be written as H<sub>2</sub>CO<sub>3</sub>.

12.




### Page 275, Quick Check

- Concentrated acids have a high molarity (ex. 6M acid) whereas a strong acid is one that ionizes completely (ex. HCl). You can have a concentrated weak acid – such as 6 M HF.
- HNO<sub>3</sub> is a strong acid so ionizes completely:  

$$\text{HNO}_3(aq) + \text{H}_2\text{O}(l) \rightarrow \text{H}_3\text{O}^+(aq) + \text{NO}_3^-(aq)$$

1.0 M      1.0 M



Because both acids are the same concentration, but HNO<sub>3</sub> is strong and CH<sub>3</sub>COOH is weak, the HNO<sub>3</sub> will have more ions in solution.

- In 0.1 M NaOH: a bright light    In 0.1 M NH<sub>3</sub>: very dim light or no light visible  

$$\text{NaOH}(aq) \rightarrow \text{Na}^+(aq) + \text{OH}^-(aq) \quad \text{NH}_3(aq) + \text{H}_2\text{O}(l) \leftrightarrow \text{NH}_4^+(aq) + \text{OH}^-(aq)$$
- No, Cl<sup>-</sup> will not accept an H<sup>+</sup> ion. HCl is a strong acid and completely ionizes.

### Page 276, Quick Check

- HF(aq) + H<sub>2</sub>O(l) ↔ H<sub>3</sub>O<sup>+</sup>(aq) + F<sup>-</sup>(aq) K<sub>a</sub> = [H<sub>3</sub>O<sup>+</sup>][F<sup>-</sup>] / [HF]
- NaOH is a strong base, which means that it ionizes completely. There is no equilibrium present, so no K<sub>b</sub>.
- CH<sub>3</sub>CH<sub>2</sub>NH<sub>2</sub>(aq) + H<sub>2</sub>O(l) ↔ OH<sup>-</sup>(aq) + CH<sub>3</sub>CH<sub>2</sub>NH<sub>3</sub><sup>+</sup>(aq)  
 K<sub>b</sub> = [OH<sup>-</sup>][CH<sub>3</sub>CH<sub>2</sub>NH<sub>3</sub><sup>+</sup>] / [CH<sub>3</sub>CH<sub>2</sub>NH<sub>2</sub>]
- Acid: HC<sub>2</sub>O<sub>4</sub><sup>-</sup>(aq) + H<sub>2</sub>O(l) ↔ H<sub>3</sub>O<sup>+</sup>(aq) + C<sub>2</sub>O<sub>4</sub><sup>2-</sup>(aq)  
 K<sub>a</sub> = [H<sub>3</sub>O<sup>+</sup>][C<sub>2</sub>O<sub>4</sub><sup>2-</sup>] / [HC<sub>2</sub>O<sub>4</sub><sup>-</sup>]  
 Base: HC<sub>2</sub>O<sub>4</sub><sup>-</sup>(aq) + H<sub>2</sub>O(l) ↔ OH<sup>-</sup>(aq) + H<sub>2</sub>C<sub>2</sub>O<sub>4</sub>(aq)  
 K<sub>b</sub> = [OH<sup>-</sup>][H<sub>2</sub>C<sub>2</sub>O<sub>4</sub>] / [HC<sub>2</sub>O<sub>4</sub><sup>-</sup>]

**Page 278, Quick Check**

- strong acid
  - strong base
  - weak base
  - weak acid
  - weak base
  - weak acid
- weak base:  $\text{NH}_3(aq) + \text{H}_2\text{O}(l) \leftrightarrow \text{NH}_4^+(aq) + \text{OH}^-(aq)$   
 $K_b = [\text{NH}_4^+][\text{OH}^-] / [\text{NH}_3]$
  - weak acid:  $\text{C}_6\text{H}_5\text{COOH}(aq) + \text{H}_2\text{O}(l) \leftrightarrow \text{H}_3\text{O}^+(aq) + \text{C}_6\text{H}_5\text{COO}^-(aq)$   
 $K_a = [\text{H}_3\text{O}^+][\text{C}_6\text{H}_5\text{COO}^-] / [\text{C}_6\text{H}_5\text{COOH}]$
  - weak base:  $\text{CN}^-(aq) + \text{H}_2\text{O}(l) \leftrightarrow \text{HCN}(aq) + \text{OH}^-(aq)$   
 $K_b = [\text{HCN}][\text{OH}^-] / [\text{CN}^-]$
  - weak acid:  $\text{HNO}_2(aq) + \text{H}_2\text{O}(l) \leftrightarrow \text{H}_3\text{O}^+(aq) + \text{NO}_2^-(aq)$   
 $K_a = [\text{H}_3\text{O}^+][\text{NO}_2^-] / [\text{HNO}_2]$
- Methanoic acid > carbonic acid > boric acid > phenol  
 $\text{HCOOH} > \text{H}_2\text{CO}_3 > \text{H}_3\text{BO}_3 > \text{C}_6\text{H}_5\text{OH}$

**Page 281, Quick Check**

- There are more oxygen atoms (which are highly electronegative) attached to the central nitrogen atom - therefore, the O-H bond in Acid 1 is more polarized than in Acid 2 (as 3 oxygen atoms pull electrons away from hydrogen to a greater extent than 2 oxygen atoms do) and hydrogen will be donated more easily in Acid 1. As a result, Acid 1 is stronger than Acid 2.
- Since the presence of more hydrogen atoms bound to the central atom strengthens the bond between hydrogen and that atom, HBr, HCl, HI > H<sub>2</sub>Se, H<sub>2</sub>S, H<sub>2</sub>O > CH<sub>4</sub> in terms of acid strength.  
 I > Br > Cl in terms of size. And as the halogen atoms' size increases, the bond length between the halogen atom and hydrogen increases, so the bond strength decreases, allowing H to be donated more easily and making the acid stronger. Therefore, HI > HBr > HCl in terms of acid strength.  
 For the same reason, Se > S > O in terms of size and H<sub>2</sub>Se > H<sub>2</sub>S > H<sub>2</sub>O in terms of acidity. Therefore, HI > HBr > HCl > H<sub>2</sub>Se > H<sub>2</sub>S > H<sub>2</sub>O > CH<sub>4</sub> in terms of acid strength.
- The more oxygen atoms are attached to the central atom, the more polarized the O-H bond is. Therefore, H<sub>2</sub>SO<sub>4</sub>, HClO<sub>4</sub> > HClO<sub>3</sub>, HBrO<sub>3</sub>, H<sub>2</sub>SO<sub>3</sub> in terms of acid strength.

Between H<sub>2</sub>SO<sub>4</sub> and HClO<sub>4</sub>, since Cl is more electronegative than H<sub>2</sub>SO<sub>4</sub> and pulls electrons away from H more than S does, hydrogen in HClO<sub>4</sub> is donated more easily than in H<sub>2</sub>SO<sub>4</sub>. Therefore, HClO<sub>4</sub> > H<sub>2</sub>SO<sub>4</sub> in terms of acid strength.

For similar reasons, H<sub>2</sub>SO<sub>3</sub> is weaker than HClO<sub>3</sub> and HBrO<sub>3</sub> (since Cl and Br are both electronegative halogens while S is not). Finally, since Br is less electronegative than Cl, the O-H bond is more polarized in chloric acid than in bromic. This allows hydrogen to be donated more easily. Therefore, HBrO<sub>3</sub> > HClO<sub>3</sub> > H<sub>2</sub>SO<sub>3</sub>.

Therefore overall, HClO<sub>4</sub> > H<sub>2</sub>SO<sub>4</sub> > HBrO<sub>3</sub> > HClO<sub>3</sub> > H<sub>2</sub>SO<sub>3</sub>.

- Methanoic acid and benzoic acid have main carbon chain lengths of 1 while ethanoic acid has a carbon chain of length 2. Therefore, ethanoic acid's O-H bond is the least polarizable. Furthermore, benzoic acid has a huge group of carbon atoms - a benzene ring (phenyl group) - attached to the 1 carbon chain that allows any charge to disperse between the many carbon & hydrogen atoms in the benzene group, so its O-H bond is the less polarizable than that of methanoic acid.  
 Therefore, methanoic acid > benzoic acid > ethanoic acid in terms of polarity of O-H bond and consequently strength.

**Page 283, Practice Problems 5.2.1**

- $\text{H}_2\text{O}_2 + \text{HSO}_3^- \leftrightarrow \text{H}_2\text{SO}_3 + \text{HO}_2^-$ ; reactants favored
  - $\text{H}_3\text{C}_6\text{H}_5\text{O}_7 + \text{NH}_3 \leftrightarrow \text{H}_2\text{C}_6\text{H}_5\text{O}_7^- + \text{NH}_4^+$ ; products favored
  - $\text{HCO}_3^- + \text{H}_2\text{PO}_4^- \leftrightarrow \text{H}_2\text{CO}_3 + \text{HPO}_4^{2-}$ ; reactants favored
- $\text{H}_3\text{AsO}_4 + \text{SO}_4^{2-} \leftrightarrow \text{H}_2\text{AsO}_4^- + \text{HSO}_4^-$ ; HSO<sub>4</sub><sup>-</sup> is stronger
  - $\text{SO}_3^{2-} + \text{Cr}(\text{H}_2\text{O})_6^{3+} \leftrightarrow \text{HSO}_3^- + \text{Cr}(\text{H}_2\text{O})_5\text{OH}^{2+}$   
 products favored so  $K_{eq} > 1$

**Page 283, 5.2 Review Questions**

- strong base
  - weak acid
  - Strong acid
  - weak base
- $\text{H}_3\text{BO}_3 + \text{H}_2\text{O} \leftrightarrow \text{H}_2\text{BO}_3^- + \text{H}_3\text{O}^+ \quad K_a = \frac{[\text{H}_2\text{BO}_3^-][\text{H}_3\text{O}^+]}{[\text{H}_3\text{BO}_3]}$   
 $\text{PO}_4^{3-} + \text{H}_2\text{O} \leftrightarrow \text{HPO}_4^{2-} + \text{OH}^- \quad K_b = \frac{[\text{HPO}_4^{2-}][\text{OH}^-]}{[\text{PO}_4^{3-}]}$

3. Both acids have the same concentration. Oxalic acid is a weak acid and hydriodic acid is strong. Conductivity depends on ions in solution. Since hydriodic acid is a strong acid, it will ionize completely. Oxalic acid is weak, so it ionizes to a lesser extent, forming fewer ions in solution.

4.  $\text{HNO}_3$  is a strong acid so ionizes completely:  
 $\text{HNO}_3(aq) + \text{H}_2\text{O}(l) \rightarrow \text{H}_3\text{O}^+(aq) + \text{NO}_3^-(aq)$   
 2.0 M      2.0 M              total [ion] = 4.0 M  
 $\text{HNO}_2$  is a weak acid, so does not ionize completely.

5. a) 6 M  $\text{NH}_3$

b) 0.001 M HCl

6. a) water < carbonic acid < citric acid < sulphurous acid < sulphuric acid

b) carbonate > ammonia > monohydrogen phosphate > fluoride > nitrite > water

7.  $\text{HC}_6\text{H}_5\text{O}_7^{2-} + \text{H}_2\text{O} \rightleftharpoons \text{C}_6\text{H}_5\text{O}_7^{3-} + \text{H}_3\text{O}^+$   
 $K_a = [\text{C}_6\text{H}_5\text{O}_7^{3-}][\text{H}_3\text{O}^+] / [\text{HC}_6\text{H}_5\text{O}_7^{2-}]$

$\text{H}_2\text{C}_6\text{H}_5\text{O}_7^- + \text{H}_2\text{O} \rightleftharpoons \text{HC}_6\text{H}_5\text{O}_7^{2-} + \text{H}_3\text{O}^+$   
 $K_a = [\text{HC}_6\text{H}_5\text{O}_7^{2-}][\text{H}_3\text{O}^+] / [\text{H}_2\text{C}_6\text{H}_5\text{O}_7^-]$

$\text{Al}(\text{H}_2\text{O})_6^{3+} + \text{H}_2\text{O} \rightleftharpoons \text{Al}(\text{H}_2\text{O})_5\text{OH}^{2+} + \text{H}_3\text{O}^+$   
 $K_a = [\text{Al}(\text{H}_2\text{O})_5\text{OH}^{2+}][\text{H}_3\text{O}^+] / [\text{Al}(\text{H}_2\text{O})_6^{3+}]$

$\text{H}_2\text{O}_2 + \text{H}_2\text{O} \rightleftharpoons \text{HO}_2^- + \text{H}_3\text{O}^+$   
 $K_a = [\text{HO}_2^-][\text{H}_3\text{O}^+] / [\text{H}_2\text{O}_2]$

8.  $\text{NH}_3 + \text{H}_2\text{O} \rightleftharpoons \text{NH}_4^+ + \text{OH}^-$

$K_b = [\text{NH}_4^+][\text{OH}^-] / [\text{NH}_3]$

$\text{C}_6\text{H}_5\text{COO}^- + \text{H}_2\text{O} \rightleftharpoons \text{C}_6\text{H}_5\text{COOH} + \text{OH}^-$   
 $K_b = [\text{C}_6\text{H}_5\text{COOH}][\text{OH}^-] / [\text{C}_6\text{H}_5\text{COO}^-]$

$\text{CH}_3\text{COO}^- + \text{H}_2\text{O} \rightleftharpoons \text{CH}_3\text{COOH} + \text{OH}^-$   
 $K_b = [\text{CH}_3\text{COOH}][\text{OH}^-] / [\text{CH}_3\text{COO}^-]$

$\text{HC}_6\text{H}_5\text{O}_7^{2-} + \text{H}_2\text{O} \rightleftharpoons \text{C}_6\text{H}_5\text{O}_7^- + \text{OH}^-$   
 $K_b = [\text{C}_6\text{H}_5\text{O}_7^-][\text{OH}^-] / [\text{HC}_6\text{H}_5\text{O}_7^{2-}]$

$\text{C}_5\text{H}_5\text{N} + \text{H}_2\text{O} \rightleftharpoons \text{C}_5\text{H}_5\text{NH}^+ + \text{OH}^-$   
 $K_b = [\text{C}_5\text{H}_5\text{NH}^+][\text{OH}^-] / [\text{C}_5\text{H}_5\text{N}]$

9.

a)  $\text{Fe}(\text{H}_2\text{O})_6^{3+}(aq) + \text{HO}_2^-(aq) \rightleftharpoons \text{H}_2\text{O}_2 + \text{Fe}(\text{H}_2\text{O})_5\text{OH}^{2+}$  products




b)  $\text{H}_2\text{SO}_3(aq) + \text{IO}_3^-(aq) \rightleftharpoons \text{HSO}_3^- + \text{HIO}_3$  reactants

c)  $\text{CN}^-(aq) + \text{H}_2\text{PO}_4^-(aq) \rightleftharpoons \text{HCN} + \text{HPO}_4^{2-}$  products

10.  $\text{H}_2\text{C}_2\text{O}_4 + \text{HCO}_3^- \rightleftharpoons \text{H}_2\text{CO}_3 + \text{HC}_2\text{O}_4^-$

11.  $\text{HClO} > \text{HBrO} > \text{H}_2\text{SiO}_3$

12. All are strong acids so completely ionize in solution to form  $\text{H}_3\text{O}^+$ :



### Page 290, Quick Check

1. energy +  $2\text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{OH}^-$

If temperature increases, equilibrium shifts right and the concentrations of  $\text{H}_3\text{O}^+$  and  $\text{OH}^-$  both increase.

$$K_w = [\text{H}_3\text{O}^+][\text{OH}^-]$$

Therefore,  $K_w$  will increase as well.

2.  $[\text{H}_3\text{O}^+] = [\text{OH}^-]$

3.  $5.4 \times 10^{-8} \text{ M} = [\text{H}_3\text{O}^+] = [\text{OH}^-]$

### Page 291, Practice Problems 5.3.1

1.  $[\text{H}_3\text{O}^+] = 0.15 \text{ M}$  because  $\text{HClO}_4$  is a strong acid;  $[\text{OH}^-] = 6.67 \times 10^{-14} \text{ M}$

acidic because  $[\text{H}_3\text{O}^+] > [\text{OH}^-]$

2.  $[\text{OH}^-] = 2s = 2.2 \times 10^{-4} \text{ M}$ ;  $[\text{H}_3\text{O}^+] = 4.5 \times 10^{-11} \text{ M}$   
 basic because  $[\text{H}_3\text{O}^+] < [\text{OH}^-]$

3.  $[\text{OH}^-] = 0.142 \text{ M}$ ;  $[\text{H}_3\text{O}^+] = 7.04 \times 10^{-14} \text{ M}$   
 basic because  $[\text{H}_3\text{O}^+] < [\text{OH}^-]$

### Page 292, Practice Problems 5.3.2

1.  $[\text{OH}^-] = 2 \times 10^{-13} \text{ M}$

2. 2 g

3.  $[\text{OH}^-] = 5.32 \times 10^{-14} \text{ M}$

### Page 294, 5.3 Review Questions

1.  $\text{NH}_3 + \text{NH}_3 \rightleftharpoons \text{NH}_4^+ + \text{NH}_2^-$

2.

$[\text{H}_3\text{O}^+]$	$[\text{OH}^-]$	Acidic, basic or neutral?
$1.7 \times 10^{-15}$	6.0M	basic
$3.2 \times 10^{-4} \text{ M}$	$3.1 \times 10^{-11}$	acidic
$1.1 \times 10^{-3}$	$9.2 \times 10^{-12} \text{ M}$	acidic
2.5M	$4.0 \times 10^{-15}$	acidic
$2.1 \times 10^{-10}$	$4.7 \times 10^{-5} \text{ M}$	basic

3.  $57.1 \text{ kJ} + 2\text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{OH}^-$

As temperature increases, equilibrium shifts right and  $[\text{H}_3\text{O}^+]$  and  $[\text{OH}^-]$  increase, so  $K_w$  increases.

4.  $[\text{OH}^-] = 5.0 \times 10^{-15} \text{ M}$


5.  $[\text{OH}^-] = 1.6 \times 10^{-8} \text{ M}$   
 acidic because  $[\text{H}_3\text{O}^+] > [\text{OH}^-]$

6. Complete the table:

Temp	$K_w$	$[\text{H}_3\text{O}^+]$	$[\text{OH}^-]$
50° C	$5.5 \times 10^{-14}$	$2.3 \times 10^{-7} \text{ M}$	$2.3 \times 10^{-7} \text{ M}$
100° C	$5.1 \times 10^{-13}$	$7.1 \times 10^{-7} \text{ M}$	$7.1 \times 10^{-7} \text{ M}$

Both are neutral at given temperatures

7.



$$K_w = 7.9 \times 10^{-15}$$

8.  $[\text{OH}^-] = 2s = 0.0211 \text{ M}$   
 $[\text{H}_3\text{O}^+] = 4.7 \times 10^{-13} \text{ M}$
9.  $[\text{OH}^-] = 4.3 \times 10^{-14} \text{ M}$
10. 3.2 g

#### Page 298, Quick Check

1.

Solution	$[\text{H}_3\text{O}^+]$	pH
orange juice	$3.2 \times 10^{-4} \text{ M}$	3.49
milk of magnesia	$2.52 \times 10^{-11} \text{ M}$	10.598
stomach acid	0.031 M	1.51

#### Page 302, Practice Problems 5.4.2

1.

Solution	$[\text{H}_3\text{O}^+]$	$[\text{OH}^-]$	pH	pOH	Acidic/Basic/Neutral?
Orange Juice	$3 \times 10^{-4} \text{ M}$	$3 \times 10^{-11} \text{ M}$	3.5	10.5	Acidic
Tears	$3.98 \times 10^{-8} \text{ M}$	$2.51 \times 10^{-7} \text{ M}$	7.40	6.60	Basic
Blood	$4.0 \times 10^{-8} \text{ M}$	$2.5 \times 10^{-7} \text{ M}$	7.40	6.60	Basic
Milk	$3.16 \times 10^{-7} \text{ M}$	$3.16 \times 10^{-8} \text{ M}$	6.500	7.500	Acidic

2.

- a) As a solution becomes more acidic, both  $[\text{H}_3\text{O}^+]$  and pOH increase and both  $[\text{OH}^-]$  and pH decrease.
  - b) A basic solution has a pOH value which is less than 7, and  $[\text{H}_3\text{O}^+]$  which is greater than  $10^{-7} \text{ M}$ .
  - c) If the pH of a solution equals 14.0, the  $[\text{OH}^-]$  equals 1 M. (1 sig. digit)
  - d) If the pOH of a solution decreases by 5, then the  $[\text{H}_3\text{O}^+]$  has decreased by a factor of  $10^5$ .
3. We would expect the  $K_w$  for water at 10°C to be less than  $10^{-14}$  because the autoionization of water is endothermic. Regardless of the temperature, pure water is always neutral and so:  
*For pure water at any temperature:*  $[\text{H}_3\text{O}^+] = [\text{OH}^-]$  and  $\text{pH} = \text{pOH}$   
 Therefore, because  $\text{pH} + \text{pOH} = \text{p}K_w$ , when we calculate the value of  $\text{p}K_w$ , the pH (and also the pOH) must be *half the value of*  $\text{p}K_w$ .  
 $\text{p}K_w = -\log(2.55 \times 10^{-15}) = 14.593$  Therefore,  $\text{pH}$  (and  $\text{pOH}$ ) =  $14.5934 = 7.297$

2. The  $[\text{H}_3\text{O}^+]$  increased because the pH decreased. A 5 unit pH decrease corresponds to a  $10^5$  times increase in hydronium concentration.
3. Initial pH (prior to dilution) = 1.00  
 After the dilution, the  $[\text{H}_3\text{O}^+] = 2.00$

#### Page 299, Practice Problems 5.4.1

1.

$[\text{H}_3\text{O}^+]$	No. of sig. digits	pH
$5.00 \times 10^{-5} \text{ M}$	3	4.301
$4.6 \times 10^{-4} \text{ M}$	2	3.34
$6.4 \times 10^{-11} \text{ M}$	2	10.19
$8.81 \times 10^{-7} \text{ M}$	3	6.055
0.00345 M	3	2.462

2. Just before the equivalence point was reached:  $[\text{H}_3\text{O}^+] = 4.5 \times 10^{-5} \text{ M}$   
 Just after the equivalence point was reached:  $[\text{H}_3\text{O}^+] = 2.2 \times 10^{-10} \text{ M}$
3. Because pH is defined as a negative logarithm, converting a negative pH value to a concentration involves the following:  
 $[\text{H}_3\text{O}^+] = 16 \text{ M}$   
 The pH scale is not necessary to conveniently express concentrations this large.

**Page 304, Practice Problems 5.4.3**

- 0.86
- 0.82
- 2.53
- 0.0513 g HCl

**Page 308, 5.4 Review Questions**

1. The pH of a solution is defined as the negative logarithm of the concentration of hydronium ions.

Equation:  $\text{pH} = -\log [\text{H}_3\text{O}^+]$

The pOH of a solution is defined as the negative logarithm of the concentration of hydroxide ions.

Equation:  $\text{pOH} = -\log [\text{OH}^-]$

2. Sorenson's logarithmic pH and pOH scales are a convenient and compact way of expressing the typically very small concentrations of hydronium and hydroxide ions respectively and the extent by which they can change in aqueous solutions

3.

$[\text{H}_3\text{O}^+]$	pH	Acidic/Basic/Neutral
$3.50 \times 10^{-6} M$	5.456	Acidic
$3.1 \times 10^{-12} M$	11.51	Basic
0.00550 M	2.260	Acidic
1.0 M	0.00	Acidic
$6.8 \times 10^{-9} M$	8.17	Basic

4.

$[\text{OH}^-]$	pOH	Acidic/Basic/Neutral
$7.2 \times 10^{-9} M$	8.14	Acidic
$2.8 \times 10^{-10} M$	9.55	Acidic
$4.88 \times 10^{-4} M$	3.312	Basic
$1.0 \times 10^{-14} M$	14.00	Acidic
0.000625 M	3.204	Basic

5.

- As a solution's pOH value and  $[\text{H}_3\text{O}^+]$  both decrease, the solution becomes more basic.
- As a solution's pH value and  $[\text{OH}^-]$  both decrease, the solution becomes more acidic.
- The product of the  $[\text{H}_3\text{O}^+]$  and  $[\text{OH}^-]$  equals  $K_w$ .
- The sum of pH and pOH equals  $\text{p}K_w$ .

6.

$[\text{H}_3\text{O}^+]$	pOH	Acidic/Basic/Neutral
0.0342 M	12.534	Acidic
$2.51 \times 10^{-6} M$	8.400	Acidic
$7.2 \times 10^{-12} M$	2.86	Basic
$1.64 \times 10^{-11} M$	3.215	Basic

7. Pure water is always neutral at any temperature and so  $\text{pH} = \text{pOH}$ .

Therefore pH 6.51 (2 sig. digits)

8.  $\text{pH} = 13.78$

9. 13.000

10. 0.81 g

11.

$[\text{H}_3\text{O}^+]$	$[\text{OH}^-]$	pOH	pH	Acidic/Basic/Neutral
$5.620 \times 10^{-5} M$	$1.779 \times 10^{-10} M$	9.7497	4.2503	Acidic
$2.22 \times 10^{-11} M$	0.000450 M	3.347	10.653	Basic
$3.2 \times 10^{-2} M$	$3.2 \times 10^{-13} M$	12.50	1.50	Acidic
$3 \times 10^{-11} M$	$3 \times 10^{-4} M$	3.5	10.5	Basic

12. 0.79

13. 11.8

14. 12.7
15. 12.24

An alternative approach would be to calculate the total number of moles of hydronium ions from the acid solutions and then subtract that total from the moles of hydroxide ions from the basic solution. The excess moles of hydroxide ions present in the final 100.0 mL are then used to calculate final pH.

16. 0.483 g HCl
17. 0.128 g LiOH

#### Page 314, Quick Check

1. The amount of water that a weak acid will convert to hydronium ions is insignificant compared to the magnitude of water's concentration (55.6 M). The concentration of water is thus assumed to be constant and so is not included in an ICE table.
2. Because the  $K_a$  of a weak acid is normally so much greater than  $K_w$ , we can assume that the initial  $[\text{H}_3\text{O}^+]$  resulting from the autoionization of water is insignificant compared to the equilibrium  $[\text{H}_3\text{O}^+]$  resulting from the ionization of this weak acid.
3. The value of  $K_a$  is small enough compared to the initial concentration of the acid that the percent of the acid which actually ionizes will not significantly change that original concentration. Remember that this assumption is only valid if the percent ionization of the weak acid is  $\leq 5\%$ .

#### Page 315, Practice Problems 5.5.1

1. 2.02
2. 2.41 (2 sig. digits)
3. 5.4 %

The simplifying assumption would not be valid in this case because the percent ionization is greater than 5 %

#### Page 316, Practice Problems 5.5.2

1. 0.017 M
2. 1.6 M
3. 0.45 M

#### Page 317, Practice Problems 5.5.3

1.  $1.0 \times 10^{-5}$
2.  $1.4 \times 10^{-4}$
3.  $6.56 \times 10^{-5}$  (The acid is most likely benzoic acid)

#### Page 319, Quick Check

1.

Weak Base	Conjugate Acid	Appropriate $K_a$ Value	$pK_a$ Value	Calculated $K_b$	$pK_b$ Value
$\text{HC}_2\text{O}_4^-$	$\text{H}_2\text{C}_2\text{O}_4$	$5.9 \times 10^{-2}$	1.23	$1.7 \times 10^{-13}$	12.77
$\text{H}_2\text{PO}_4^-$	$\text{H}_3\text{PO}_4$	$7.5 \times 10^{-3}$	2.12	$1.3 \times 10^{-12}$	11.88
$\text{HPO}_4^{2-}$	$\text{H}_2\text{PO}_4^-$	$6.2 \times 10^{-8}$	7.21	$1.6 \times 10^{-7}$	6.79
$\text{NO}_2^-$	$\text{HNO}_2$	$4.6 \times 10^{-4}$	3.34	$2.2 \times 10^{-11}$	10.66
$\text{HC}_6\text{H}_5\text{O}_7^{2-}$	$\text{H}_2\text{C}_6\text{H}_5\text{O}_7^-$	$1.7 \times 10^{-5}$	4.77	$5.9 \times 10^{-10}$	9.23
$\text{HCO}_3^-$	$\text{H}_2\text{CO}_3$	$4.3 \times 10^{-7}$	6.37	$2.3 \times 10^{-8}$	7.63
$\text{CN}^-$	$\text{HCN}$	$4.9 \times 10^{-10}$	9.31	$2.0 \times 10^{-5}$	4.69

#### Page 321, Practice Problems 5.5.4

1. 11.05
2.  $x = [\text{OH}^-] = 5.8 \times 10^{-6} \text{ M}$  so  $\text{pOH} = -\log(5.8 \times 10^{-6}) = 5.24$   
 $\text{pH} = 14.00 - 5.24 = 8.76$  and  $[\text{H}_3\text{O}^+] = 10^{-8.76} = 1.7 \times 10^{-9} \text{ M}$

3. (a) Methylammonium,  $\text{CH}_3\text{NH}_3^+$   
 $K_a = 2.3 \times 10^{-11}$   
 $pK_a = 10.64$   
 (b) Hydrazinium,  $\text{N}_2\text{H}_5^+$   
 $K_a = 5.9 \times 10^{-9}$   
 $pK_a = 8.23$

**Page 322, Practice Problems 5.5.5**

- 0.87 g  $\text{C}_2\text{H}_5\text{NH}_2$
- 0.49 M
- $6.0 \times 10^{-3}$  M

**Page 324, Practice Problems 5.5.6**

- $K_b = 0.020$ ; Percent ionization = 20. %  
 This calculated  $K_b$  value is greater than the  $K_b$  given in the sample problem ( $4.4 \times 10^{-4}$ ). This indicates that the temperature of the solution is higher than  $25^\circ\text{C}$ .
- $K_b = 2 \times 10^{-6}$  (1 sig. digits)  
 $pK_b = -\log(1.6 \times 10^{-6}) = 5.79$
- $3.3 \times 10^{-6}$

**Page 328, 5.5 Review Questions**

- $\text{pH} = -\log(0.01342) = 1.87$  (2 sig. digits)     $[\text{H}_3\text{O}^+] = 10^{-1.87} = 0.013$  M  
 $\text{pOH} = 14.00 - 1.87 = 12.13$      $[\text{OH}^-] = 10^{-12.13} = 7.4 \times 10^{-13}$  M
- $\text{pH} = 4.02$ ; Percent ionization = 0.095 %
- $\text{pH} = 2.873$ ;  $K_a = 1.80 \times 10^{-5}$  (The unknown acid is acetic acid.)
- 8.9 g HF
- 0.16 M  $\text{H}_3\text{PO}_4$
- 0.11 g  $\text{H}_2\text{C}_2\text{O}_4$
- $2.9 \times 10^{-8}$
- $4.9 \times 10^{-5}$
- 

Conjugate Acid	Conjugate Base	$K_a$ for acid	$pK_a$	$K_b$ for base	$pK_b$
$\text{HNO}_2$	$\text{NO}_2^-$	$4.6 \times 10^{-4}$	3.34	$2.2 \times 10^{-11}$	10.66
$\text{H}_2\text{O}_2$	$\text{HO}_2^-$	$2.4 \times 10^{-12}$	11.62	$4.2 \times 10^{-3}$	2.38
$\text{C}_6\text{H}_5\text{OH}$	$\text{C}_6\text{H}_5\text{O}^-$	$1.3 \times 10^{-10}$	9.89	$7.7 \times 10^{-5}$	4.11
$\text{HSO}_4^-$	$\text{SO}_4^{2-}$	$1.2 \times 10^{-2}$	1.92	$8.3 \times 10^{-13}$	12.08

10.

Conjugate Acid	Conjugate Base	$K_a$ for acid	$pK_a$	$K_b$ for base	$pK_b$
$\text{H}_2\text{PO}_4^-$	$\text{HPO}_4^{2-}$	$6.2 \times 10^{-8}$	7.21	$1.6 \times 10^{-7}$	6.79
$\text{H}_2\text{C}_6\text{H}_5\text{O}_7^-$	$\text{HC}_6\text{H}_5\text{O}_7^{2-}$	$1.7 \times 10^{-5}$	4.77	$5.9 \times 10^{-10}$	9.23
$\text{H}_3\text{BO}_3$	$\text{H}_2\text{BO}_3^-$	$7.3 \times 10^{-10}$	9.14	$1.4 \times 10^{-5}$	4.86
$\text{HCO}_3^-$	$\text{CO}_3^{2-}$	$5.6 \times 10^{-11}$	10.25	$1.8 \times 10^{-4}$	3.75

- 11.39
- 12.23
- 0.10 M
- 9.1 L
- $4.2 \times 10^{-4}$